

ALCAM Antibody / Activated leukocyte cell adhesion molecule [clone rALCAM/9811] (V5884)

Catalog No.	Formulation	Size
V5884-100UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	100 ug
V5884-20UG	0.2 mg/ml in 1X PBS with 0.05% BSA, 0.05% sodium azide	20 ug
V5884SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **MOUSE MONOCLONAL**

Bulk quote request

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2b, kappa
Clone Name	rALCAM/9811
UniProt	Q13740
Localization	Cell membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml
Limitations	This ALCAM/Activated leukocyte cell adhesion molecule antibody is available for research use only.

Application Notes

Optimal dilution of the ALCAM/Activated leukocyte cell adhesion molecule antibody should be determined by the researcher.

Immunogen

Recombinant full-length human CD166 protein was used as the immuonogen for the ALCAM/Activated leukocyte cell adhesion molecule antibody.

Storage

ALCAM/Activated leukocyte cell adhesion molecule antibody with sodium azide - store at 2 to 8oC; antibody without sodium azide - store at -20 to -80oC.

