


## Alkyl-DHAP synthase Antibody / AGPS (RQ7215)


| Catalog No. | Formulation                                           | Size   |
|-------------|-------------------------------------------------------|--------|
| RQ7215      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**


|                           |                                                                                                                                                                           |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Availability</b>       | 1-3 business days                                                                                                                                                         |
| <b>Species Reactivity</b> | Human                                                                                                                                                                     |
| <b>Format</b>             | Antigen affinity purified                                                                                                                                                 |
| <b>Host</b>               | Rabbit                                                                                                                                                                    |
| <b>Clonality</b>          | Polyclonal (rabbit origin)                                                                                                                                                |
| <b>Isotype</b>            | Rabbit IgG                                                                                                                                                                |
| <b>Purity</b>             | Antigen affinity purified                                                                                                                                                 |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2% Trehalose                                                                                                                                 |
| <b>UniProt</b>            | O00116                                                                                                                                                                    |
| <b>Localization</b>       | Cytoplasmic (peroxisome)                                                                                                                                                  |
| <b>Applications</b>       | Western Blot : 0.5-1ug/ml<br>Immunohistochemistry (FFPE) : 2-5ug/ml<br>Immunofluorescence : 5ug/ml<br>Flow Cytometry : 1-3ug/million cells<br>Direct ELISA : 0.1-0.5ug/ml |
| <b>Limitations</b>        | This Alkyl-DHAP synthase antibody is available for research use only.                                                                                                     |




IHC staining of FFPE human liver cancer tissue with Alkyl-DHAP synthase antibody.  
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human adenocarcinoma of lung tissue with Alkyl-DHAP synthase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human adenocarcinoma of lung tissue with Alkyl-DHAP synthase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human esophageal squamous cell carcinoma tissue with Alkyl-DHAP synthase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human cervical cancer tissue with Alkyl-DHAP synthase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human tonsil tissue with Alkyl-DHAP synthase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human SiHa cells with Alkyl-DHAP synthase antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) HeLa, 2) K562, 3) HepG2, 4) RT4, 5) HEL, 6) Caco-2 and 7) SiHa cell lysate with Alkyl-DHAP synthase antibody. Predicted molecular weight ~73 kDa.



Flow cytometry testing of human U937 cells with Alkyl-DHAP synthase antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Alkyl-DHAP synthase antibody.

## Description

This gene is a member of the FAD-binding oxidoreductase/transferase type 4 family. It encodes a protein that catalyzes the second step of ether lipid biosynthesis in which acyl-dihydroxyacetonephosphate (DHAP) is converted to alkyl-DHAP by the addition of a long chain alcohol and the removal of a long-chain acid anion. The protein is localized to the inner aspect of the peroxisomal membrane and requires FAD as a cofactor. Mutations in this gene have been associated with rhizomelic chondrodysplasia punctata, type 3 and Zellweger syndrome.

## Application Notes

Optimal dilution of the Alkyl-DHAP synthase antibody should be determined by the researcher.

## Immunogen

Recombinant human protein (amino acids D154-L658) was used as the immunogen for the Alkyl-DHAP synthase antibody.

## Storage

After reconstitution, the Alkyl-DHAP synthase antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.