


## Alpha 2 macroglobulin receptor Antibody / CD91 / LRP1 [clone LRP1/6784] (V4267)

Catalog No.	Formulation	Size
V4267-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4267-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4267SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	LRP1/6784
Purity	Protein A/G affinity
UniProt	Q07954
Localization	Cytoplasm, Nucleus, Cell surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This Alpha 2 macroglobulin receptor antibody is available for research use only.



IHC staining of FFPE human lung tissue with Alpha 2 macroglobulin receptor antibody (clone LRP1/6784) Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

### Description

Members of the LDL receptor gene family, including LDLR (low density lipoprotein receptor), LEP1 (low density

lipoprotein related protein), megalin (also designated GP330), VLDLR (very low density lipoprotein receptor) and apoER2 are characterized by a cluster of cysteine-rich class A repeats, epidermal growth factor (EGF)-like repeats, YWTD repeats and an O-linked sugar domain. LRP1, also designated LRP and Alpha-2-Macroglobulin receptor, is an endocytic receptor that mediates the uptake of at least 15 ligands, including Alpha-2-Macroglobulin and apoE. LRP1 is cleaved into a membrane subunit and an extracellular subunit, which remain non-covalently associated. Proper folding and trafficking of LRP1 is facilitated by the receptor-associated protein (RAP), a molecular chaperone. The uptake of all known ligands through LRP1 can be blocked by RAP, which induces a conformational change in the receptor that renders it unable to bind ligands. LRP1, which is expressed in brain, liver and lung, is also implicated in Alzheimer's disease (AD), as the human LRP gene localizes to a potential AD locus on chromosome 12.

## Application Notes

Optimal dilution of the Alpha 2 macroglobulin receptor antibody should be determined by the researcher.

## Immunogen

A recombinant fragment of human protein was used as the immunogen for the Alpha 2 macroglobulin receptor antibody.

## Storage

Aliquot the Alpha 2 macroglobulin receptor antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.