


## Alpha Tubulin Antibody [clone TUBA/3038] (V8404)

Catalog No.	Formulation	Size
V8404-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8404-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8404SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	TUBA/3038
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P68363
<b>Localization</b>	Cell surface, Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This Alpha Tubulin antibody is available for research use only.




IHC staining of FFPE human lung with Alpha Tubulin antibody (clone TUBA/3038).  
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

#### Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Alpha Tubulin antibody (clone TUBA/3038). These results demonstrate the foremost specificity of the TUBA/3038 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free Alpha Tubulin antibody (clone TUBA/3038) as confirmation of integrity and purity.

## Description

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulin. This MAAb recognizes an epitope of alpha-tubulin. The alpha-tubulin is globular protein that exists in cells as part of soluble alpha/beta-tubulin dimer or it is polymerized into microtubules. In different species it is coded by multiple tubulin genes that form tubulin classes (in human 6 genes). Expressed tubulin genes are named tubulin isotypes. Some of the tubulin isotypes are expressed ubiquitously, while some have more restricted tissue expression. Alpha-tubulin is also subject of numerous post-translational modifications. Tubulin isotypes and their posttranslational modifications are responsible for multiple tubulin charge variants - tubulin isoforms. Heterogeneity of alpha-tubulin is concentrated in C-terminal structural domain

## Application Notes

Optimal dilution of the Alpha Tubulin antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 1-115 from the human protein was used as the immunogen for the Alpha Tubulin antibody.

## Storage

Store the Alpha Tubulin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).