


# **Androgen Receptor Antibody [clone ANDRC-1] (V7101)**

Catalog No.	Formulation	Size
V7101-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7101-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7101SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7101IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	ANDRC-1
Purity	Protein G affinity chromatography
UniProt	P10275
Localization	Nuclear
Applications	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1) Prediluted IHC Only Format: incubate for 30 min at RT (2)
Limitations	This Androgen Receptor antibody is available for research use only.



IHC testing of FFPE human prostate carcinoma and Androgen Receptor antibody (clone ANDRC-1).

#### **Description**

The Androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. [RefSeq]

#### **Application Notes**

Optimal dilution of the Androgen Receptor antibody should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

### **Immunogen**

Recombinant full-length human AR protein was used as the immunogen for the Androgen Receptor antibody.

#### **Storage**

Store the Androgen Receptor antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).