


Anti-Androgen Receptor Antibody [clone SPM335] (V2639)

Catalog No.	Formulation	Size
V2639-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2639-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2639SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2639IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM335
Purity	Protein G affinity chromatography
UniProt	P10275
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This anti-Androgen Receptor antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human prostate carcinoma stained with anti-Androgen Receptor antibody (SPM335).

Description

Recognizes a protein of 100kDa, which is identified as androgen receptor (AR). It reacts with full length, and the newly described A form of the receptor. It does not cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. well differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marker of hormone-responsiveness and thus it may be useful in identifying patients likely to benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This mAb is superb for staining of formalin/paraffin tissues.

Application Notes

Optimal dilution of the anti-Androgen Receptor antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 302-318 (STEDTAEYSPFKGGYTK) from human AR were used as the immunogen for the anti-Androgen Receptor antibody.

Storage

Store the anti-Androgen Receptor antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).