


# **Anti-EGFR Antibody (F49183)**


| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F49183-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F49183-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

## **Bulk quote request**


| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Antigen affinity purified  |
| Clonality          | Polyclonal (rabbit origin)   |
| Isotype            | Rabbit Ig  |
| Purity             | Antigen affinity   |
| UniProt            | P00533   |
| Applications       | Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Flow Cytometry : 1:10-1:50 |
| Limitations        | This anti-EGFR antibody is available for research use only.                  |



Western blot analysis of lysate from A431, HUVEC, PC3 cell line (left to right) using anti-EGFR antibody diluted at 1:1000 for each lane.



Anti-EGFR antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human placenta tissue.



Anti-EGFR antibody flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

### **Description**

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer.

#### **Application Notes**

Titration of the anti-EGFR antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 1004-1033 from the human protein was used as the immunogen for this anti-EGFR antibody.

#### **Storage**

Aliquot the anti-EGFR antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.