


## Anti-Glycophorin A Antibody [clone SPM599] (V2563)


Catalog No.	Formulation	Size
V2563-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2563-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2563SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2563IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	SPM599
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P02724
<b>Localization</b>	Cytoplasmic, membranous
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 1-2ug/ml
<b>Limitations</b>	This anti-Glycophorin A antibody is available for research use only.



IHC testing of FFPE human placenta stained with anti-Glycophorin A antibody (clone SPM599). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



IHC testing of FFPE human angiosarcoma stained with anti-Glycophorin A antibody (clone SPM599). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



Western blot testing of human K562 cell lysate with anti-Glycophorin A antibody. Expected molecular weight: ~19 kDa (glycosylated monomer), ~38 kDa (glycosylated dimer).

## Description

Recognizes a sialoglycoprotein of 39kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. Glycophorins A (GPA) and B (GPB), which are single, trans-membrane sialoglycoproteins. GPA is the carrier of blood group M and N specificities, while GPB accounts for S and U specificities. GPA and GPB provide the cells with a large mucin like surface and it has been suggested this provides a barrier to cell fusion, so minimizing aggregation between red blood cells in the circulation.

## Application Notes

Optimal dilution of the anti-Glycophorin A antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Recombinant human GPA protein was used as the immunogen for the anti-Glycophorin A antibody.

## Storage

Store the anti-Glycophorin A antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).