
# Anti Mullerian Hormone Antibody / AMH [clone AMH/7354] (V4593)


| Catalog No. | Formulation | Size |
|---|---|---|
| V4593-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4593-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4593SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Purified |
| **Host** | Mouse |
| **Clonality** | Monoclonal (mouse origin) |
| **Isotype** | Mouse IgG1, kappa |
| **Clone Name** | AMH/7354 |
| **Purity** | Protein A/G affinity |
| **UniProt** | P03971 |
| **Localization** | Secreted |
| **Applications** | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| **Limitations** | This Anti Mullerian Hormone antibody is available for research use only. |



IHC staining of FFPE human testis tissue with Anti-Mullerian Hormone antibody (clone AMH/7354). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




SDS-PAGE analysis of purified, BSA-free Anti-Mullerian Hormone antibody (clone AMH/7354) as confirmation of integrity and purity.

## Description

The transforming growth factor beta (TGF beta) superfamily is composed of numerous growth and differentiation factors, including TGF beta1-3, Mullerian inhibiting substance (MIS), growth/differentiation factor (GDF) 1-9, bone morphogenic protein (BMP) 2-8, glial cell line-derived neurotrophic factor (GDNF), Inhibin Alpha, beta-A, beta-B and beta-C, Lefty and Nodal. Members of the TGF beta superfamily are involved in embryonic development and adult tissue homeostasis. The MIS glycoprotein is produced by the Sertoli cells of the testes. Fetal testes produce both MIS and testosterone, the presence of which result in male offspring. Absence of MIS and testosterone in a developing fetus results in the induction of Mullerian duct differentiation, and Wolffian duct development is not induced. Testosterone induces the differentiation of the Wolffian ducts whereas MIS causes regression of the Muellerian duct. MIS inhibits the growth of tumors derived from tissues of Mullerian duct origin. MIS can also inhibit the autophosphorylation of the EGF receptor in vitro. Defects in anti-Muellerian hormone are the cause of persistent Muellerian duct syndrome type I (PMDS-1). PMDS-1 is a form of male pseudo hermaphroditism characterized by a failure of Muellerian duct regression in otherwise normal males.

## Application Notes

Optimal dilution of the Anti Mullerian Hormone antibody should be determined by the researcher.

## Immunogen

A recombinant fragment of the C-terminus of the human protein (within amino acids 460-560) was used as the immunogen for the Anti Mullerian Hormone antibody.

## Storage

Aliquot the Anti Mullerian Hormone antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.