
# Apolipoprotein A I Antibody (R31886)


| Catalog No. | Formulation | Size |
|---|---|---|
| R31886 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**


| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Rat |
| **Format** | Antigen affinity purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit IgG |
| **Purity** | Antigen affinity |
| **Buffer** | Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide |
| **UniProt** | P04639 |
| **Localization** | Cytoplasmic, membranous |
| **Applications** | Western Blot : 0.1-0.5ug/ml<br>IHC (FFPE) : 0.5-1ug/ml |
| **Limitations** | This Apolipoprotein A I antibody is available for research use only. |




Western blot testing of 1) rat liver and 2) rat testis lysate with Apolipoprotein A I antibody. Predicted molecular weight ~30 kDa.



IHC testing of FFPE rat liver with Apolipoprotein A I antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat kidney with Apolipoprotein A I antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

## Description

Apolipoprotein A-1, also known as APOA1, is a human protein with a specific role in lipid metabolism. It binds to lipopolysaccharide or endotoxin, and has a major role in the anti-endotoxin function of HDL. The gene is mapped to 11q23. And it is a single polypeptide chain with 243 amino acid residues of known primary amino acid sequence. The ApoA-I protein promotes cholesterol efflux from tissues to the liver for excretion. It is a cofactor for lecithin cholesterolacyltransferase (LCAT) which is responsible for the formation of most plasma cholesteryl esters. ApoA-I is also isolated as a prostacyclin (PGI2) stabilizing factor, and thus may have an anticlotting effect. Defects in the gene encoding it are associated with HDL deficiencies, including Tangier disease, and with systemic non-neuropathic amyloidosis. Additionally, ApoA-I overexpression promotes macrophage-specific reverse cholesterol transport.

For a broadly validated polyclonal option, see our APOA1 Antibody / Apolipoprotein A1 Antibody page.

## Application Notes

Optimal dilution of the Apolipoprotein A I antibody should be determined by the researcher.

## Immunogen

Amino acids 25-259 of rat Apolipoprotein A I were used as the immunogen for the Apolipoprotein A I antibody.

## Storage

After reconstitution, the Apolipoprotein A I antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.


Copyright © NSJ Bioreagents. All rights reserved.