
APRT Antibody (RQ4891)


Catalog No.	Formulation	Size
RQ4891	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P08030
Applications	Western blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 1-2ug/ml
Limitations	This APRT antibody is available for research use only.



Western blot testing of human 1) T-47D, 2) HEK293, 3) K562 and 4) Caco-2 lysate with APRT antibody at 0.5ug/ml. Predicted molecular weight ~20 kDa.



IHC staining of FFPE human breast cancer with APRT antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.

Description

Adenine phosphoribosyltransferase (APRTase) is an enzyme encoded by the APRT gene, found in humans on chromosome 16. It belongs to the purine/pyrimidine phosphoribosyltransferase family. A conserved feature of this gene is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis. Two transcript variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the APRT antibody should be determined by the researcher.

Immunogen

Amino acids RAEVVECVSLVELTSLKGRERL from the mouse protein were used as the immunogen for the APRT antibody.

Storage

After reconstitution, the APRT antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.