


AR Antibody / Nuclear receptor subfamily 3 group C member 4 [clone MSVA-367R] (V6092)

Catalog No.	Formulation	Size
V6092-100UG	Antibody in 1X PBS with 0.05% BSA, 0.05% sodium azide	100 ug
V6092-20UG	Antibody in 1X PBS with 0.05% BSA, 0.05% sodium azide	20 ug

Recombinant RABBIT MONOCLONAL

Bulk quote request

Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MSVA-367R
UniProt	P10275
Localization	Cytoplasm, Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This AR/Nuclear receptor subfamily 3 group C member 4 antibody is available for research use only.



Androgen receptor Rabbit Recombinant Monoclonal Antibody (MSVA-367R) tested on many normal and cancer tissues. The immunohistochemistry staining in these tissues aligns with the expression data in Human Protein Atlas.

Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121oC in pH 7.8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:150 at 37oC for 60 minutes. Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.

Application Notes

- Optimal dilution of the AR/Nuclear receptor subfamily 3 group C member 4 antibody should be determined by the researcher.
- This AR/Nuclear receptor subfamily 3 group C member 4 antibody is recombinantly produced by expression in CHO

cells.

Immunogen

Synthetic peptide corresponding to residues (within amino acids 1-100) of Androgen Receptor was used as the immunogen for the AR/Nuclear receptor subfamily 3 group C member 4 antibody.

Storage

AR/Nuclear receptor subfamily 3 group C member 4 antibody with sodium azide - store at 2 to 8oC; antibody without sodium azide - store at -20 to -80oC.