


ARG1 Antibody (F51459)

Catalog No.	Formulation	Size
F51459-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F51459-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P05089
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Flow Cytometry : 1:10-1:50
Limitations	This ARG1 antibody is available for research use only.

95 72 55 -	Western blot analysis of ARG1 antibody and MDA-MB231 lysate. Predicted molecular weight ~35kDa.
43 34 - ◀	
26 -	



IHC analysis of FFPE human hepatocarcinoma stained with ARG1 antibody



ARG1 antibody flow cytometric analysis of MDA-MB231 cells (green) compared to a <u>negative control</u> (blue). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

Description

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exist (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type I isoform encoded by this gene, is a cytosolic enzyme and expressed predominantly in the liver as a component of the urea cycle. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia.

Application Notes

Titration of the ARG1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 293-322 from the human protein was used as the immunogen for this ARG1 antibody.

Storage

Aliquot the ARG1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.