


ARG1 Antibody / Liver Arginase [clone 2B12] (RQ5861)


Catalog No.	Formulation	Size
RQ5861	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat, Monkey
Format	Antigen affinity purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	2B12
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P05089
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This ARG1 antibody is available for research use only.



Flow cytometry testing of human Jurkat cells with ARG1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ARG1 antibody.



Flow cytometry testing of human SiHa cells with ARG1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ARG1 antibody.



Western blot testing of 1) rat liver, 2) mouse liver and 3) monkey liver lysate with ARG1 antibody. Predicted molecular weight ~35 kDa.

Description

ARG1 (arginase, live) is a cytosolic enzyme and expressed predominantly in the liver as a component of the urea cycle. The isoform encoded by ARG1, referred to as the liver, or A-I, isoform, contributes 98% of the arginase activity in liver but is also present in red cells. Using a rat liver ARG1 cDNA clone to probe a human liver cDNA library, Haraguchi et al. (1987) isolated and characterized a cDNA corresponding to the ARG1 gene. The ARG1 gene is mapped on 6q23.2 and the arginase gene contains 8 exons. By immunologic studies, 90% of the arginase in red blood cell and liver was precipitated by the antibody, whereas only 50% of the arginase in kidney, brain, and the gastrointestinal tract reacted with it. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia. Two transcript variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the ARG1 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids E25-D183) was used as the immunogen for the ARG1 antibody.

Storage

After reconstitution, the ARG1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.