


## ARK1C1 Antibody / Aldo-keto Reductase Family 1 Member C1 [clone AKR1C1/9070] (V5584)


Catalog No.	Formulation	Size
V5584-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5584-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5584SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat, Rabbit, Hamster, Guinea pig
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	AKR1C1/9070
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q04828
<b>Localization</b>	Cytoplasm
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml (Human/Mouse/Rat/Hamster/Rabbit/Guinea pig)
<b>Limitations</b>	This ARK1C1 antibody is available for research use only.




Western blot testing of human liver tissue lysate using AKR1C1 antibody (clone AKR1C1/9070). Predicted molecular weight ~37 kDa.



IHC staining of FFPE human hepatocellular carcinoma tissue with AKR1C1 antibody (clone AKR1C1/9070). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free AKR1C1 antibody (clone AKR1C1/9070) as confirmation of integrity and purity.



AKR1C1 Antibody Multi-Species Liver Panel WB. Western blot analysis of (1) human liver, (2) mouse liver, (3) rat liver, (4) hamster liver, (5) guinea pig liver, and (6) HepG2 cell lysates using AKR1C1 Antibody (clone AKR1C1/9070) detects a band at approximately 37 kDa, consistent with the predicted molecular weight of AKR1C1 / Aldo-keto reductase family 1 member C1. The conserved banding pattern across species and in HepG2 cells supports reliable detection of AKR1C1 in hepatic tissue, aligning with its role in steroid metabolism and detoxification pathways.

## Description

DDH / AKR1C1 is a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reaction of progesterone to the inactive form 20-alpha-hydroxy-progesterone.

This antibody can be compared with our [AKR1C1 Antibody](#) (clone AKR1C1/9063) for consistent detection of AKR1C1 across steroid metabolism and detoxification studies.

## Application Notes

Optimal dilution of the AKR1C1 antibody should be determined by the researcher.

## Immunogen

Recombinant human full-length AKR1C1 protein was used as the immunogen for the AKR1C1 antibody.

## Storage

Aliquot the AKR1C1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

