


## ARK1C1 Antibody / Aldo-keto Reductase Family 1 Member C1 [clone AKR1C1/9314] (V5586)


Catalog No.	Formulation	Size
V5586-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5586-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5586SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	AKR1C1/9314
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q04828
<b>Localization</b>	Cytoplasm
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This ARK1C1 antibody is available for research use only.



IHC staining of FFPE human hepatocellular carcinoma tissue with AKR1C1 antibody (clone AKR1C1/9314). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free AKR1C1 antibody (clone AKR1C1/9314) as confirmation of integrity and purity.

## Description

DDH / AKR1C1 is a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reaction of progesterone to the inactive form 20-alpha-hydroxy-progesterone.

## Application Notes

Optimal dilution of the AKR1C1 antibody should be determined by the researcher.

## Immunogen

Recombinant human full-length AKR1C1 protein was used as the immunogen for the AKR1C1 antibody.

## Storage

Aliquot the AKR1C1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.