


ARK1C1 Antibody / Aldo-keto Reductase Family 1 Member C1 [clone AKR1C1/9319] (V5588)


Catalog No.	Formulation	Size
V5588-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5588-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5588SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	AKR1C1/9319
Purity	Protein A/G affinity
UniProt	Q04828
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml
Limitations	This ARK1C1 antibody is available for research use only.



Western blot testing of human HepG2 cell lysate using AKR1C1 antibody (clone AKR1C1/9319). Predicted molecular weight ~37 kDa.



IHC staining of FFPE human hepatocellular carcinoma tissue with AKR1C1 antibody (clone AKR1C1/9319). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free AKR1C1 antibody (clone AKR1C1/9319) as confirmation of integrity and purity.

Description

DDH / AKR1C1 is a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reaction of progesterone to the inactive form 20-alpha-hydroxy-progesterone.

Application Notes

Optimal dilution of the ARK1C1 antibody should be determined by the researcher.

Immunogen

Recombinant human full-length AKR1C1 protein was used as the immunogen for the ARK1C1 antibody.

Storage

Aliquot the ARK1C1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.