


## ATF2 Antibody / Activating Transcription Factor 2 [clone IA-1] (RQ5447)

Catalog No.	Formulation	Size
RQ5447	Antibody in PBS with 0.02% sodium azide, 50% glycerol and 0.4-0.5mg/ml BSA	100 ul

[Bulk quote request](#)

Availability	1-2 weeks
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	IA-1
Purity	Affinity purified
UniProt	P15336
Applications	Western Blot : 1:500-1:2000
Limitations	This ATF2 antibody is available for research use only.



Western blot testing of human HeLa cell lysate with ATF2 antibody. Expected molecular weight: 65-70 kDa.

### Description

The ATF2 gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription. This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. The encoded protein may also be involved in cell's

DNA damage response independent of its role in transcriptional regulation. [RefSeq]

## Application Notes

Optimal dilution of the ATF2 antibody should be determined by the researcher.

## Immunogen

A synthetic peptide specific to human Activating Transcription Factor 2 was used as the immunogen for the ATF2 antibody.

## Storage

Store the ATF2 antibody at -20oC.