


## ATF6 Antibody [clone AF6229] (N1111)

Catalog No.	Formulation	Size
N1111-100UG	0.5 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
N1111-25UG	0.5 mg/ml antibody in PBS with 0.1 mg/ml BSA (US source) and 0.05% sodium azide	25 ug

[Bulk quote request](#)

<b>Availability</b>	2-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	IgG1, kappa
<b>Clone Name</b>	AF6229
<b>Purity</b>	Protein G purified monoclonal antibody
<b>Buffer</b>	1X PBS, pH 7.4
<b>Gene ID</b>	22926
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 2-4ug/ml (2) Immunohistochemistry (FFPE) : 5ug/ml (1)
<b>Limitations</b>	This ATF6 antibody is available for research use only.



Western blot testing of human samples with ATF6 antibody (AF6229) at 2ug/ml.  
Predicted molecular weight: 75~90kDa.

## Description

ER stress can cause the accumulation of unfolded proteins and lead to the activation of the UPR (unfolded protein response). Cells have developed an intracellular signaling pathway to deal with this stress, and Activating Transcription Factor 6 (ATF6) is a major transducer of the UPR. Upon activation, ATF6 separates from GRP78 and is cleaved. The ~50 kD cytosolic portion of ATF6 translocates to the nucleus, transmitting the ER stress signal. In the nucleus, ATF6 activates the transcription of UPR-responsive genes by binding to their ERSE (ER stress response element) region.

## Application Notes

ATF6 is a 670 amino acid protein that migrates in western blot between 70~90 kD due to glycosylation.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.

## Immunogen

A recombinant protein fragment from the N-terminal region of human ATF6 was used as the immunogen for this antibody.

## Storage

Aliquot the antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

## Alternate Names

Activating Transcription Factor 6