


ATG5 Antibody [clone ATG5/2492] (V7623)


| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V7623-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V7623-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V7623SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request


| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1, kappa |
| Clone Name | ATG5/2492 |
| Purity | Protein G affinity chromatography |
| UniProt | Q9H1Y0 |
| Localization | Cytoplasmic |
| Applications | ELISA (order BSA-free Format For Coating) : Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This ATG5 antibody is available for research use only. |




IHC staining of FFPE human colon carcinoma with ATG5 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.




IHC staining of FFPE human endometrial carcinoma with ATG5 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.




IHC staining of FFPE human ovarian carcinoma with ATG5 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



IHC staining of FFPE human brain with ATG5 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



Western blot testing of human ThP1 and brain lysate with ATG5 antibody. Predicted molecular weight ATG5: ~32 kDa; ATG5/ATG12 heterodimer: ~56 kDa.



SDS-PAGE analysis of purified, BSA-free ATG5 antibody as confirmation of integrity and purity.

Description

The protein encoded by this gene, in combination with autophagy protein 12, functions as an E1-like activating enzyme in a ubiquitin-like conjugating system. The encoded protein is involved in several cellular processes, including autophagic vesicle formation, mitochondrial quality control after oxidative damage, negative regulation of the innate antiviral immune response, lymphocyte development and proliferation, MHC II antigen presentation, adipocyte differentiation, and apoptosis. The ATG5 protein is essential for autophagy; a process that is usually beneficial for cells to self-degrade their own components when they are no longer useful.

Application Notes

Optimal dilution of the ATG5 antibody should be determined by the researcher.

Immunogen

A recombinant human partial protein (amino acids 1-119) was used as the immunogen for the ATG5 antibody.

Storage

Store the ATG5 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).