


Basigin Antibody / CD147 / Emmprin [clone BSG/7952] (V4597)


| Catalog No. | Formulation | Size |
|----------------|-------------------------------------------------------------------------|--------|
| V4597-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4597-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4597SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request


| | |
|---------------------------|------------------------------------------------------------------------------------|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG |
| Clone Name | BSG/7952 |
| Purity | Protein A/G affinity |
| UniProt | P35613 |
| Localization | Cell surface |
| Applications | Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| Limitations | This Basigin antibody is available for research use only. |




Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using Basigin antibody (clone BSG/7952). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to be specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.




Western blot testing of human THP-1 cell lysate with Basigin antibody (clone BSG/7952).
Expected molecular weight: 27-66 kDa depending on level of glycosylation.



IHC staining of FFPE human placental tissue with Basigin antibody (clone BSG/7952) at 2ug/ml. Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human renal cell carcinoma tissue with Basigin antibody (clone BSG/7952) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Basigin antibody (clone BSG/7952) as confirmation of integrity and purity.

Description

Basigin is expressed more intensely on thymocytes than on mature peripheral blood T cells. It is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. It stimulates the production of interstitial collagenase, gelatinase A, stromelysin-1 and various metalloproteinases (MMPs) by fibroblasts. These enzymes are important factors in cancer invasion and metastasis.

Application Notes

Optimal dilution of the Basigin antibody should be determined by the researcher.

Immunogen

Recombinant full-length human protein was used as the immunogen for the Basigin antibody.

Storage

Aliquot the Basigin antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

