


BCAT1 Antibody / Branched-chain Aminotransferase 1 (R30615)


Catalog No.	Formulation	Size
R30615	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P54687
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This BCAT1 antibody is available for research use only.



IHC staining of FFPE human pancreatic cancer tissue with BCAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human ovarian adenoma tissue with BCAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) Raji and 2) MOLT4 cell lysate with BCAT1 antibody.
Predicted molecular weight ~43 kDa.

Description

Branched-chain Aminotransferase 1 is also known as BCT1. The gene is highly expressed early in embryogenesis, and during organogenesis its expression is localized to the neural tube, the somites, and the mesonephric tubules. The gene is also expressed in several MYC-based tumors. Lack of the enzyme BCAT1 can cause auxotroph, a kind of auxotrophic mutant in Chinese-hamster ovary cells that lacks the ability to grow if alpha-ketoisovaleric acid, alpha-ketoisocaproic acid, and alpha-keto-beta-methylvaleric acid are substituted for valine, leucine, and isoleucine in the culture medium.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the BCAT1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the C-terminus of human BCAT1 (KLASRILSKLTDIQYGR) was used as the immunogen for this BCAT1 antibody.

Storage

After reconstitution, the BCAT1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.