
BCL2L1 Antibody / Bcl-XL [clone 31B78] (FY13325)

Catalog No.	Formulation	Size
FY13325	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA	100 ul


Recombinant RABBIT MONOCLONAL

Bulk quote request


Availability	2-3 weeks
Species Reactivity	Human, Mouse, Rat
Format	Liquid
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	31B78
Purity	Affinity chromatography
Buffer	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
UniProt	Q07817
Applications	Western blot: 1:500-1:2000 Immunohistochemistry: 1:50-1:200 Immunoprecipitation: 1:50
Limitations	This BCL2L1 antibody is available for research use only.




BCL2L1 Antibody WB. Western blot analysis of human HeLa, Jurkat, HepG2, MCF7, and SH-SY5Y cell lysates using BCL2L1 Antibody demonstrates a prominent immunoreactive band at approximately 26-32 kDa, consistent with expression of BCL2L1/Bcl-XL. Bcl-XL is the major anti-apoptotic isoform encoded by the BCL2L1 gene and functions as a key regulator of mitochondrial apoptosis by inhibiting pro-apoptotic BCL2 family proteins and promoting cellular survival. A faint band near 50 kDa is observed in some cell lines and may represent an SDS-resistant Bcl-XL dimer or higher-order protein complex. This higher molecular weight species is substantially less abundant than the predominant monomeric form. Antibody incubation was performed at 1:1000 dilution for 1 hour at room temperature. Predicted molecular weight of Bcl-XL: approximately 26-30 kDa.



BCL2L1 Antibody Hematopoietic Cell WB. Western blot analysis of human Ramos (lane 1) and K562 (lane 2) cell lysates using BCL2L1 Antibody demonstrates distinct immunoreactive bands at approximately 26-32 kDa, consistent with expression of Bcl-XL, the major anti-apoptotic isoform encoded by the BCL2L1 gene. Bcl-XL functions as a critical regulator of mitochondrial apoptosis by inhibiting activation of pro-apoptotic BCL2 family proteins and promoting cellular survival under conditions of physiological and cellular stress. The observed banding pattern is consistent with endogenous BCL2L1 expression in hematopoietic-derived cell lines and supports specific detection of the protein by western blot analysis. Predicted molecular weight: approximately 26 kDa. Commonly observed molecular weight: 26-32 kDa.



BCL2L1 Antibody Cross-Species WB. Western blot analysis of mouse heart, mouse liver, mouse spleen, mouse lung, mouse skin, rat heart, and rat liver tissue lysates using BCL2L1 Antibody demonstrates a predominant immunoreactive band at approximately 26-32 kDa, consistent with expression of Bcl-XL, the major anti-apoptotic isoform encoded by the BCL2L1 gene. Bcl-XL functions as a critical regulator of mitochondrial apoptosis by inhibiting pro-apoptotic BCL2 family proteins and promoting cellular survival across diverse tissues. A faint band near 50 kDa is observed in selected samples and may represent an SDS-resistant Bcl-XL dimer or higher-order protein complex. This higher molecular weight species is substantially less abundant than the predominant monomeric form. Antibody incubation was performed at 1:1000 dilution for 1 hour at room temperature. Predicted molecular weight: approximately 26 kDa. Commonly observed molecular weight: 26-32 kDa.



BCL2L1 Antibody Brain Tissue WB. Western blot analysis of mouse brain and rat brain tissue lysates using BCL2L1 Antibody demonstrates prominent immunoreactive bands at approximately 26-32 kDa, consistent with expression of Bcl-XL, the major anti-apoptotic isoform encoded by the BCL2L1 gene. Bcl-XL is highly expressed in neuronal tissues where it contributes to mitochondrial integrity, neuronal survival, and protection against apoptosis-induced cell death. The observed banding pattern supports conserved BCL2L1 expression across rodent brain tissues and is consistent with the established role of Bcl-XL in maintaining survival of post-mitotic neuronal populations. Antibody incubation was performed at 1:1000 dilution for 1 hour at room temperature. Predicted molecular weight: approximately 26 kDa. Commonly observed molecular weight: 26-32 kDa.

Description

BCL2L1 antibody detects Bcl-x, encoded by the BCL2L1 gene. Bcl-x is a member of the Bcl-2 protein family that regulates mitochondrial outer membrane permeability and controls apoptosis. This protein is best known for existing in multiple isoforms with opposing functions: the larger isoform Bcl-xL acts as an anti-apoptotic factor, while the shorter isoform Bcl-xS promotes apoptosis. BCL2L1 antibody provides researchers with an essential reagent to investigate cell survival, stress responses, and oncogenesis.

Bcl-x functions by interacting with other members of the Bcl-2 family, including Bax, Bak, and Bad. Research using BCL2L1 antibody has shown that the anti-apoptotic isoform Bcl-xL sequesters pro-apoptotic factors, preventing them from permeabilizing the mitochondrial outer membrane. This blocks cytochrome c release and subsequent caspase activation, preserving cell survival under stress conditions. Conversely, Bcl-xS antagonizes Bcl-xL and promotes apoptosis by facilitating mitochondrial permeabilization. The ratio of isoform expression therefore determines whether a cell undergoes survival or programmed cell death.

Studies with BCL2L1 antibody have revealed important roles for Bcl-x in development. Bcl-xL is required for the survival of immature neurons and erythroid progenitors during embryogenesis. Knockout models demonstrate embryonic lethality when Bcl-x is absent, highlighting its essential role in protecting developing tissues from apoptosis. These findings

underscore the protein's fundamental role in balancing proliferation with programmed cell death during development.

In cancer research, Bcl-x has emerged as a major regulator of tumor cell survival and therapeutic resistance. Research using BCL2L1 antibody has shown that many cancers, including lymphomas, leukemias, lung carcinoma, and breast carcinoma, overexpress Bcl-xL. This overexpression prevents apoptosis induced by chemotherapy or radiation, leading to poor therapeutic outcomes. Small molecule inhibitors targeting Bcl-xL are under active development as strategies to re-sensitize cancer cells to apoptosis. The presence of high Bcl-xL levels is often associated with worse prognosis, demonstrating its clinical relevance.

Beyond oncology, Bcl-x is also critical in cardiovascular and neurodegenerative disease. Studies with BCL2L1 antibody have shown that Bcl-xL protects cardiomyocytes during ischemia-reperfusion injury and reduces apoptosis in neurons exposed to oxidative or excitotoxic stress. However, prolonged survival of damaged cells may contribute to pathological remodeling in the heart or persistence of dysfunctional neurons in neurodegenerative disorders. This dual role highlights the complexity of targeting Bcl-x in disease therapy.

Immune regulation also relies on Bcl-x. Research using BCL2L1 antibody has demonstrated that Bcl-x isoform balance shapes lymphocyte development, particularly during thymocyte maturation. Anti-apoptotic activity ensures survival of certain T-cell subsets, while regulation of Bcl-x levels contributes to immune tolerance and selection. These functions link Bcl-x to immune homeostasis as well as immune-mediated disease.

BCL2L1 antibody is widely used in western blotting, immunohistochemistry, and flow cytometry. Western blotting distinguishes between Bcl-xL and Bcl-xS isoforms, immunohistochemistry reveals tissue-specific expression patterns, and flow cytometry measures survival regulation within immune populations. These approaches make BCL2L1 antibody valuable for basic research, translational oncology, and clinical immunology.

By supplying validated BCL2L1 antibody reagents, NSJ Bioreagents supports research into apoptosis regulation, cancer biology, and stress adaptation. Detection of Bcl-x provides insight into how Bcl-2 family proteins integrate pro-survival and pro-apoptotic signals to determine cell fate across development, immunity, and disease.

Learn more about Bcl-XL function, apoptosis regulation, mitochondrial survival signaling, and BCL2L1 biology on our Bcl-XL Antibody / Anti-Apoptotic Protein Antibody page.

Application Notes

Optimal dilution of the BCL2L1 antibody should be determined by the researcher.

Immunogen

A synthesized peptide derived from human Bcl-XL was used as the immunogen for the BCL2L1 antibody.

Storage

Store the BCL2L1 antibody at -20oC.

Copyright © NSJ Bioreagents. All rights reserved.