

BCL6 Antibody / B-Cell Lymphoma 6 Protein [clone PCR-P-BCL6-1B1] (V5021)


| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V5021-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V5021-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V5021SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

[Bulk quote request](#)


| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG2b |
| Clone Name | PCR-P-BCL6-1B1 |
| Purity | Protein A/G affinity |
| UniProt | P41182 |
| Localization | Nucleus, Mitochondria |
| Applications | Flow Cytometry : 1-2ug/million cells |
| Limitations | This BCL6 antibody is available for research use only. |



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using BCL6 antibody (clone PCR-P-BCL6-1B1). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.



Flow cytometry testing of PFA-fixed human HeLa cells with BCL6 antibody (clone PCR-P-BCL6-1B1) followed by goat anti-mouse IgG-CF488 (blue); Red = unstained cells.



SDS-PAGE analysis of purified, BSA-free BCL6 antibody (clone PCR-P-BCL6-1B1) as confirmation of integrity and purity.

Description

Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large (L&H) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.

Application Notes

Optimal dilution of the BCL6 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 528-601) from the human protein was used as the immunogen for the BCL6 antibody.

Storage

Aliquot the BCL6 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.