


# Beta Amyloid Antibody / APP [clone APP/4469] (V5428)


Catalog No.	Formulation	Size
V5428-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5428-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5428SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	APP/4469
Purity	Protein A/G affinity
UniProt	P05067
Localization	Cell membrane, cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Beta Amyloid antibody is available for research use only.



IHC staining of FFPE human brain tissue with Beta Amyloid antibody (clone APP/4469). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Beta Amyloid antibody (clone APP/4469) as confirmation of integrity and purity.

## **Description**

Proteolytic cleavage of the Amyloid protein precursor (APP) gives rise to the b-Amyloid and Amyloid A4 proteins, which are present in human platelets. Amyloid deposition is associated with type II diabetes, Down syndrome and a variety of neurological disorders, including Alzheimer's disease. The Amyloid precursor protein (APP) undergoes alternative splicing, resulting in several isoforms. Proteolytic cleavage of APP leads to the formation of the 4 kDa Amyloid b/A4 Amyloid protein. This protein is involved in the formation of neurofibrillary tangles and plaques that characterize the senile plaques of Alzheimer's patients. APLP1 (Amyloid precursor-like protein 1) and APLP2 are structurally similar to APP. Human APLP2 is a membrane-bound sperm protein that contains a region highly homologous to the transmembrane-cytoplasmic domains of APP found in brain plaques of Alzheimer's disease patients.

### **Application Notes**

Optimal dilution of the Beta Amyloid antibody should be determined by the researcher.

#### **Immunogen**

A portion of amino acids 1-100 of the amyloid beta protein was used as the immunogen for the Beta Amyloid antibody. This antibody recognizes an epitope in the region of amino acids 8-16 of the amyloid precursor protein.

#### **Storage**

Aliquot the Beta Amyloid antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.