


## Beta Catenin Antibody (C-Terminal Region) [clone 691CT11.2.1] (F54099)

Catalog No.	Formulation	Size
F54099-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54099-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat, Monkey
<b>Predicted Reactivity</b>	Bovine
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1
<b>Clone Name</b>	691CT11.2.1
<b>Purity</b>	Protein G purified
<b>UniProt</b>	P35222
<b>Applications</b>	Western Blot : 1:4000
<b>Limitations</b>	This Beta Catenin antibody is available for research use only.



Western blot testing of Beta Catenin antibody at 1:4000: Lane 1) human HEK293, 2) (h) HeLa, 3) monkey COS-7, 4) rat C6 and 5) mouse NIH3T3 cell lysate. Predicted molecular weight ~85 kDa, but routinely observed at 90-95 kDa.

### Description

Beta Catenin / CTNNB1 is a key downstream component of the canonical Wnt signaling pathway. In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome.

This CTNNB1 antibody complements our [Beta-Catenin Antibody / CTNNB1 Antibody \(clone CTNNB1/2030R\)](#) for broader analysis of CTNNB1 expression and localization.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the Beta Catenin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 687-721 from human Beta Catenin was used as the immunogen for the Beta Catenin antibody.

## Storage

Aliquot the Beta Catenin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.