


Biotin Antibody [clone rBTN/8819] (V5320)

Catalog No.	Formulation	Size
V5320-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5320-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5320SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	All species
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rBTN/8819
Purity	Protein A/G affinity
UniProt	Not Applicable
Applications	Flow Cytometry : 0.5-1ug/million cells Immunofluorescence : 0.5-1ug/ml Western Blot : 0.5-2ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 minutes at RT
Limitations	This Biotin antibody is available for research use only.



IHC staining of FFPE human kidney tissue with biotinylated Lambda Light Chain antibody probe followed by Biotin antibody (clone rBTN/8819).



SDS-PAGE analysis of purified, BSA-free Biotin antibody (clone rBTN/8819) as confirmation of integrity and purity.

Description

It recognizes both the free and protein-conjugated (either soluble or cell bound) form of biotin. This mAb is highly specific to biotin and shows no cross-reaction with other structurally related compounds. It has a very high affinity for biotin and is excellent for use in various amplification techniques. In some applications, localization of biotinylated probes with avidin produces unacceptably high background staining. Anti-biotin antibody may be substituted to decrease this noise.

Application Notes

Optimal dilution of the Biotin antibody should be determined by the researcher.

Immunogen

Biotinylated sheep immunoglobulin was used as the immunogen for the Biotin antibody.

Storage

Aliquot the Biotin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.