


## Bleomycin hydrolase Antibody / BLMH (F54739)


Catalog No.	Formulation	Size
F54739-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54739-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	Q13867
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 1:25 Western Blot : 1:500-1:1000
<b>Limitations</b>	This Bleomycin hydrolase antibody is available for research use only.



Western blot testing of human 1) HL60, 2) CCRF-CEM, 3) SH-SY5Y, 4) MOLT4 and 5) mouse Neuro-2a cell lysate with Bleomycin hydrolase antibody. Predicted molecular weight ~53 kDa.



Western blot testing of 1) human L562, 2) mouse stomach and 3) rat stomach lysate with Bleomycin hydrolase antibody. Predicted molecular weight ~53 kDa.



IHC testing of FFPE human skin tissue with Bleomycin hydrolase antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

## Description

The normal physiological role of BLM hydrolase is unknown, but it catalyzes the inactivation of the antitumor drug BLM (a glycopeptide) by hydrolyzing the carboxamide bond of its B-aminoalaninamide moiety thus protecting normal and malignant cells from BLM toxicity (By similarity).

## Application Notes

The stated application concentrations are suggested starting points. Titration of the Bleomycin hydrolase antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 212-242 from the human protein was used as the immunogen for the Bleomycin hydrolase antibody.

## Storage

Aliquot the Bleomycin hydrolase antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.