

BMPR1B Antibody (R31786)


Catalog No.	Formulation	Size
R31786	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	O00238
Localization	Cytoplasmic, membrane
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This BMPR1B antibody is available for research use only.



Western blot testing of human placenta lysate with BMPR1B lysate. Predicted/observed molecular weight: 55-60 kDa.



IHC testing of FFPE human prostate cancer with BMPR1B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

Description

BMPR1B (Bone Morphogenetic Protein Receptor Type IB), also known as ALK6, is a protein which in humans is encoded by the BMPR1B gene. BMPR1B is a member of the bone morphogenetic protein (BMP) receptor family of transmembrane serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta superfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transduce their signals through the formation of heteromeric complexes of 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. By analysis of a monochromosome hybrid mapping panel and by FISH, Astrom et al. (1999) mapped the BMPR1B gene to chromosome 4q22-q24. Ide et al. (1997) compared BMP receptor expression in normal and cancerous prostate tissues. While BMPR1A and BMPR2 were expressed at similar levels in all prostate tissues, BMPR1B was expressed at a significantly reduced level in cancerous prostate tissue.

Application Notes

Optimal dilution of the BMPR1B antibody should be determined by the researcher.

Immunogen

Amino acids 14-184 of human BMPR1B were used as the immunogen for the BMPR1B antibody.

Storage

After reconstitution, the BMPR1B antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.