


BrdU Antibody [clone 85-2C8] (V3112)

Catalog No.	Formulation	Size
V3112-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3112-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3112SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3112IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

 Citations (11)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	All species
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	85-2C8
Purity	Protein G affinity chromatography
UniProt	Not Applicable
Localization	Nuclear
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This BrdU antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded mouse small intestine stained with BrdU antibody (85-2C8).

Description

It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting S-phase cells, providing useful information on the aggressiveness of tumors.

Application Notes

Optimal dilution of the BrdU antibody should be determined by the researcher.

1. For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 min at RT followed by digestion with Trypsin at 1mg/ml PBS, 10 min at 37oC.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Bromodeoxyuridine conjugated to hemocyanine was used as the immunogen for the BrdU antibody.

Storage

Store the BrdU antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).