


## **BrdU Antibody [clone BRD.3] (V3110)**

Catalog No.	Formulation	Size
V3110-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3110-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3110SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3110IHC-7ML	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	7 ml

# Citations (11)

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	All species
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	BRD.3
Purity	Protein G affinity chromatography
Localization	Nuclear
Applications	Flow Cytometry: 0.5-1ug/10^6 cells Immunofluorescence: 0.5-1ug/ml Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This BrdU antibody is available for research use only.



IHC analysis of formalin-fixed, paraffin-embedded mouse small intestine stained with BrdU antibody (clone BRD.3).

It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting S-phase cells, providing useful information on the aggressiveness of tumors.

#### **Application Notes**

Optimal dilution of the BrdU antibody should be determined by the researcher.

- 1. For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 min at RT followed by digestion with Trypsin at 1mg/ml PBS, 10 min at 37oC.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

#### **Immunogen**

Bromodeoxyuridine conjugated to BSA was used as the immunogen for the BrdU antibody.

#### **Storage**

Store the BrdU antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).