


BTBD9 Antibody / BTB/POZ domain-containing protein 9 [clone BTBD9/7501] (V5123)


Catalog No.	Formulation	Size
V5123-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5123-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5123SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	BTBD9/7501
Purity	Protein A/G affinity
UniProt	Q96Q07
Localization	Nucleus, Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This BTBD9 antibody is available for research use only.



IHC staining of FFPE human fallopian tube tissue with BTBD9 antibody (clone BTBD9/7501). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free BTBD9 antibody (clone BTBD9/7501) as confirmation of integrity and purity.

Description

BTBD9 (BTB/POZ domain-containing protein 9) is a 612 amino acid protein that contains one BTB/POZ domain and one BACK (BTB/Kelch associated) domain. The BTB/POZ domain mediates homomeric and heteromeric POZPOZ interactions and is common to transcriptional regulators involved in chromatin modeling. In several BTB/POZ containing proteins, including Bcl-6 and the promyelocytic leukemia zinc-finger (PLZF) oncoprotein, this domain interacts with the SMRT/N-CoR-mSin3A HDAC complex and is directly involved in repressing and silencing gene transcription. When this domain is deleted, as with the oncogenic PLZF-RAR chimera of promyelocytic leukemias, this transcriptional repression is attenuated. This suggests that BTBD9 may play a role in transcription regulation. Genetic variations in the gene that encodes BTBD9 have been associated with susceptibility to restless legs syndrome type 6 (RLS6), a condition characterized by an uncontrollable urge to move the legs while resting.

Application Notes

Optimal dilution of the BTBD9 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 200-400) from the human protein was used as the immunogen for the BTBD9 antibody.

Storage

Aliquot the BTBD9 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.