


## BUB1 Antibody (RQ6759)


Catalog No.	Formulation	Size
RQ6759	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	O43683
<b>Localization</b>	Nuclear, cytoplasmic
<b>Applications</b>	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This BUB1 antibody is available for research use only.




IHC staining of FFPE rat thymus tissue with BUB1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse thymus tissue with BUB1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human bladder cancer tissue with BUB1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HeLa, 2) human 293T, 3) human K562 and 4) mouse testis tissue lysate with BUB1 antibody. Predicted molecular weight ~122 kDa.

## Description

BUB1, also known as mitotic checkpoint serine/threonine kinase, is an enzyme that in humans is encoded by the BUB1 gene. It is mapped to 2q13. BUB1 is first identified in genetic screens of *Saccharomyces cerevisiae*. The protein is bound to kinetochores and plays a key role in the establishment of the mitotic spindle checkpoint and chromosome congression. The mitotic checkpoint kinase is evolutionary conserved in organisms as diverse as *Saccharomyces cerevisiae* and humans. Loss-of-function mutations or absence of BUB1 has been reported to result in aneuploidy, chromosomal instability (CIN) and premature senescence. The protein kinase BUB1 possesses versatile and distinct functions during the cell cycle, mainly in the SAC and chromosome alignment during metaphase.

## Application Notes

Optimal dilution of the BUB1 antibody should be determined by the researcher.

## Immunogen

Recombinant human protein (amino acids E26-K1085) was used as the immunogen for the BUB1 antibody.

## Storage

After reconstitution, the BUB1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

