

# c-Myc Antibody [clone CT14.G4] (V3125)

Catalog No.	Formulation	Size
V3125-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3125-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3125SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CT14.G4
Purity	Protein G affinity chromatography
UniProt	P01106
Localization	Nuclear
Applications	Flow Cytometry : 0.5-1ug/10^6 cells Immunofluorescence : 1-2ug/ml
Limitations	This c-Myc antibody is available for research use only.



### **Description**

The c-Myc protein is a transcription factor, which is encoded by the c-Myc gene on human chromosome 8q24. c-Myc is

commonly activated in a variety of tumor cells and plays an important role in cellular proliferation, differentiation, apoptosis and cell cycle progression. The phosphorylation of c-Myc has been investigated and previous studies have suggested a functional association between phosphorylation at Thr58/Ser62 by glycogen synthase kinase 3, cyclin dependent kinase, ERK2 and C-Jun N terminal Kinase (JNK) in cell proliferation and cell cycle regulation. Studies also have shown that c-Myc is essential for tumor cell development in vasculogenesis and angiogenesis that distribute blood throughout the cells, and which brought extensive attention in the development of new therapeutic approach for cancer treatment.

#### **Application Notes**

The optimal dilution of the c-Myc antibody for each application should be determined by the researcher.

#### **Immunogen**

Amino acids 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from the human protein were used as the immunogen for this c-Myc antibody.

#### **Storage**

Store the c-Myc antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).