


c-Myc Antibody [clone MYC909] (V2746)

Catalog No.	Formulation	Size
V2746-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2746-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2746SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2746IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MYC909
Purity	Protein G affinity chromatography
UniProt	P01106
Localization	Nuclear
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This c-Myc antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human cervical carcinoma stained with c-Myc antibody (MYC909).

Description

It recognizes a transcription factor of 64-67kDa, identified as c-myc. This mAb shows no cross-reaction with v-myc. c-Myc is a transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes. [UniProt]

Application Notes

The stated application concentrations are suggested starting amounts. Optimal dilution of the c-Myc antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the c-Myc antibody.

Storage

Store the c-Myc antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).