


## c-Myc Antibody (R30632)

Catalog No.	Formulation	Size
R30632	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
<b>UniProt</b>	P01106
<b>Applications</b>	Western Blot : 0.5-1ug/ml
<b>Limitations</b>	This c-Myc antibody is available for research use only.



Western blot testing of c-Myc antibody and Lane 1: HeLa; 2: HEPG2 lysate

## Description

C-Myc is an oncogene that functions both in the stimulation of cell proliferation and in apoptosis. c-Myc elicits its oncogenic activity by causing immortalization, and to a lesser extent the transformation of cells, in addition to several other mechanisms. The c-MYC proto-oncogene encodes a transcription factor that is critical for cell growth and proliferation. It is one of the genes frequently altered in cancer cells in which it exhibits constitutive activity. Downregulation of c-Myc is critical for 2-Methoxyestradiol(2ME2)-induced oxidative stress and apoptosis in AML cells. And its up-regulation is important for promoting lymphocyte cell division, and demonstrating that GFP-c-Myc expression is a marker of proliferating lymphocytes in vivo.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the c-Myc antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the C-terminus of human c-Myc (RKRREQLKHKLEQLRNSCA) was used as the immunogen for this c-Myc antibody.

## Storage

After reconstitution, the c-Myc antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.