


## c-RET Antibody [clone RET/7690] (V4089)


Catalog No.	Formulation	Size
V4089-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4089-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4089SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	RET/7690
Purity	Protein A/G affinity
UniProt	P07949
Localization	Cytoplasm, Cell Surface, Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This c-RET antibody is available for research use only.



IHC staining of FFPE human parathyroid gland with RET Proto-oncogene antibody (clone RET/7690). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate carcinoma tissue with RET Proto-oncogene antibody (clone RET/7690). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

The Ret proto-oncogene is structurally related to the growing family of tyrosine kinase transmembrane receptors and is involved in GDNF signaling. RET expression is reported in several regions of the central nervous system; in the developing cranial nerve ganglia and a subset of cells within dorsal root ganglia, in motor neurons in the spinal cord and hindbrain, in neuro-retina and the growing tips of the renal collecting ducts in developing kidney. Alterations in RET gene are associated with diseases including papillary thyroid carcinoma, multiple endocrine neoplasia (type 2A and 2B), familial medullary thyroid carcinoma, and a congenital developmental disorder known as Hirschsprung's disease.

## Application Notes

Optimal dilution of the c-RET antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein (within amino acids 702-848) from the human protein was used as the immunogen for the c-RET antibody.

## Storage

Aliquot the c-RET antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.