


CAD Antibody Cocktail / CALD1 / Caldesmon (HMW) [clone CALD1/820 + h-CALD] (V2943)

Catalog No.	Formulation	Size
V2943-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2943-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2943SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2943IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CALD1/820 + h-CALD
Purity	Protein G affinity chromatography
UniProt	Q05682
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Limitations	This CAD antibody cocktail is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human uterus stained with CAD antibody (CALD1/820 + h-CALD).

Description

Two closely related variants of human caldesmon have been identified which are different in their electrophoretic mobility and cellular distribution. The h-caldesmon variant (120-150kDa) is predominantly expressed in smooth muscle whereas l-caldesmon (70-80kDa) is found in non-muscle tissue and cells. Neither of the two variants has been detected in skeletal muscle. Caldesmon is a developmentally regulated protein involved in smooth muscle and non-muscle contraction.

Application Notes

Optimal dilution of the CAD antibody cocktail should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant full-length human protein (CALD1/820) and crude human uterus extract (h-CALD) were used as the immunogen for the CAD antibody cocktail.

Storage

Store the CAD antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).