


## Cadherin 17 Antibody / CDH17 [clone CDH17/2612] (V5638)


Catalog No.	Formulation	Size
V5638-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5638-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5638SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	CDH17/2612
<b>Purity</b>	Protein G affinity
<b>UniProt</b>	Q12864
<b>Localization</b>	Cell membrane, cytoplasm
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 1-2ug/ml
<b>Limitations</b>	This Cadherin 17 antibody is available for research use only.



Western blot testing of human small intestine tissue lysate with Cadherin 17 antibody (clone CDH17/2612). Predicted molecular weight ~92 kDa but may be observed at higher molecular weights due to glycosylation.



IHC staining of FFPE human colon tissue with Cadherin 17 antibody (clone CDH17/2612). Inset: PBS used in place of primary Ab (secondary Ab negative control).  
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Cadherin 17 antibody (clone CDH17/2612) as confirmation of integrity and purity.

## Description

Cadherin 17 (also known as LI Cadherin) is part of a family of Calcium-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including beta-catenin, to regulate cadherin function. LI-cadherin (for liver-intestine-cadherin) expression is restricted to liver and intestine tissues and is specifically localized to the basolateral domain of hepatocytes and enterocytes.

## Application Notes

Optimal dilution of the Cadherin 17 antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 242-418 from human Cadherin 17 protein was used as the immunogen for the Cadherin 17 antibody.

## Storage

Aliquot the Cadherin 17 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.