


# Calbindin 2 Antibody / CALB2 / Calretinin [clone CALB2/2807] (V7544)


Catalog No.	Formulation	Size
V7544-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7544-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7544SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CALB2/2807
Purity	Protein G affinity chromatography
UniProt	P22676
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Calbindin 2 antibody is available for research use only.




IHC testing of FFPE human mesothelioma with Calbindin 2 antibody (clone CALB2/2807). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.




IHC testing of FFPE human testicular carcinoma with Calbindin 2 antibody (clone CALB2/2807). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human cerebellum with Calbindin 2 antibody (clone CALB2/2807). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.




Western blot testing of human brain lysate with Calbindin 2 antibody (clone CALB2/2807). Expected molecular weight ~29 kDa.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Calbindin 2 antibody (clone CALB2/2807). These results demonstrate the foremost specificity of the CALB2/2807 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free Calbindin 2 antibody (clone CALB2/2807) as confirmation of integrity and purity.

#### **Description**

It recognizes a protein of about 29kDa, which is identified as Calretinin (also known as Calbindin 2). Calretinin is a vitamin D-dependent calcium-binding protein involved in calcium signaling. It is present in subsets of neurons throughout the brain and spinal chord, including sensory ganglia. Antibody to calretinin is useful in differentiating mesothelioma from adenocarcinomas of the lung. It also aids in differentiating adrenal cortical neoplasms from pheochromocytomas.

#### **Application Notes**

Optimal dilution of the Calbindin 2 antibody should be determined by the researcher.

## **Immunogen**

A portion of amino acids 23-242 from the human protein was used as the immunogen for the Calretinin antibody.

## **Storage**

Store the Calbindin 2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).