


# Calreticulin Antibody / CALR (F54897)


Catalog No.	Formulation	Size
F54897-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54897-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	P27797
Localization	Cytoplasmic
Applications	Western Blot : 1:500-1:1000 Flow Cytometry : 1:10-1:50 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:10-1:50
Limitations	This Calreticulin antibody is available for research use only.



IHC testing of FFPE human colon carcinoma tissue with Calreticulin antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Western blot testing of 1) human HeLa, 2) human SH-SY5Y and 3) rat C6 cell lysate with Calreticulin antibody. Predicted molecular weight ~48 kDa but routinely observed at 55~60 kDa.



Flow cytometry testing of human HepG2 cells with Calreticulin antibody; Blue=isotype control, Green= Calreticulin antibody.

### **Description**

Calreticulin is a multifunctional protein that acts as a major Ca(2+)-binding (storage) protein in the lumen of the endoplasmic reticulum. It is also found in the nucleus, suggesting that it may have a role in transcription regulation. Calreticulin binds to the synthetic peptide KLGFFKR, which is almost identical to an amino acid sequence in the DNA-binding domain of the superfamily of nuclear receptors. Calreticulin binds to antibodies in certain sera of systemic lupus and Sjogren patients which contain anti-Ro/SSA antibodies, it is highly conserved among species, and it is located in the endoplasmic and sarcoplasmic reticulum where it may bind calcium. The amino terminus of calreticulin interacts with the DNA-binding domain of the glucocorticoid receptor and prevents the receptor from binding to its specific glucocorticoid response element. Calreticulin can inhibit the binding of androgen receptor to its hormone-responsive DNA element and can inhibit androgen receptor and retinoic acid receptor transcriptional activities in vivo, as well as retinoic acid-induced neuronal differentiation. Thus, calreticulin can act as an important modulator of the regulation of gene transcription by nuclear hormone receptors. Systemic lupus erythematosus is associated with increased autoantibody titers against calreticulin but calreticulin is not a Ro/SS-A antigen. Earlier papers referred to calreticulin as an Ro/SS-A antigen but this was later disproven. Increased autoantibody titer against human calreticulin is found in infants with complete congenital heart block of both the IgG and IgM classes.

## **Application Notes**

The stated application concentrations are suggested starting points. Titration of the Calreticulin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

Recombinant human protein was used as the immunogen for the Calreticulin antibody.

#### **Storage**

Aliquot the Calreticulin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.