


CARM1 Antibody / PRMT4 [clone CARM1/7426] (V4583)


Catalog No.	Formulation	Size
V4583-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4583-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4583SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	CARM1/7426
Purity	Protein A/G affinity
UniProt	Q86X55
Localization	Nucleus, Cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CARM1 antibody is available for research use only.



IHC staining of FFPE human prostate carcinoma tissue with CARM1 antibody (clone CARM1/7426). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate tissue with CARM1 antibody (clone CARM1/7426). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CARM1 antibody (clone CARM1/7426) as confirmation of integrity and purity.

Description

CARM1 (coactivator-associated arginine methyltransferase 1), also known as protein arginine N-methyltransferase 4 (PRMT4), is a 585 amino acid nuclear and cytoplasmic protein belonging to the protein arginine N-methyltransferase family. As a protein arginine N-methyltransferase, CARM1 is capable of catalyzing the transfer of methyl groups from S-adenosylmethionine to the guanidino group nitrogen atoms of arginine residues in certain proteins involved in mRNA stability, DNA packaging and transcriptional regulation. The methyltransferase activity of CARM1 has been found to be negatively regulated through phosphorylation at a conserved serine residue. CARM1 acts as a positive regulator for multiple transcription factors and functions as a secondary coactivator through its association with p160 coactivators. CARM1 exists as two alternatively spliced isoforms, and is encoded by a gene that maps to human chromosome 19p13.2.

Application Notes

Optimal dilution of the CARM1 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 200-400) from the human protein was used as the immunogen for the CARM1 antibody.

Storage

Aliquot the CARM1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.