
Caspase 3 Antibody [clone 15G8] (RQ4905)


Catalog No.	Formulation	Size
RQ4905	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1
Clone Name	15G8
Purity	Purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P42574
Applications	Western blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 1-2ug/ml Immunofluorescence (FFPE): 2-4ug/ml Flow cytometry: 1-3ug/million cells
Limitations	This Caspase 3 antibody is available for research use only.




Immunofluorescent staining of FFPE human MCF7 cells with Caspase 3 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IHC staining of FFPE human placenta with Caspase 3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human intestinal cancer with Caspase 3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human placenta and 2) human SW579 lysate with Caspase 3 antibody. The pro form is seen at ~32kD and active caspase-3 seen at ~17 kDa and ~12 kDa.



Flow cytometry testing of human HepG2 cells with Caspase 3 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Caspase 3 antibody.

Description

Caspase 3 is a caspase protein which interacts with Survivin, XIAP, CFLAR, Caspase 8, HCLS1, Deleted in Colorectal Cancer, TRAF3 and GroEL. This gene which is located on 4q35 encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes that undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. And the caspase-3 activation in heart failure sequentially cleaves SRF and generates a truncated SRF that appears to function as a dominant-negative transcription factor. Additionally, the caspase-3 influence on bone mineral density should be considered in any in vivo application of caspase-3 inhibitors to the treatment of human disease. In erythroid precursors undergoing terminal differentiation, Hsp70 prevents active CASP3 from cleaving GATA1 and inducing apoptosis.

Application Notes

Optimal dilution of the Caspase 3 antibody should be determined by the researcher.

Immunogen

Amino acids T67-D175 from the human protein were used as the immunogen for the Caspase 3 antibody.

Storage

After reconstitution, the Caspase 3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.