


Cathepsin D Antibody / CTSD [clone CTPD-1] (V7644)


Catalog No.	Formulation	Size
V7644-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7644-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V7644SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CTPD-1
Purity	Protein G affinity chromatography
UniProt	P07339
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml Western Blot : 1-2ug/ml
Limitations	This Cathepsin D antibody is available for research use only.



IHC staining of FFPE human liver tissue with Cathepsin D antibody (clone CTPD-1).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



Western blot testing of human liver lysate with CTSD antibody (clone CTPD-1).

Description

Cathepsin D is a ubiquitously expressed lysosomal aspartyl protease involved in the normal degradation of proteins. It is synthesized as an inactive 52kDa preprocathepsin D that is cleaved and glycosylated to form a 48kDa procathepsin D and then further cleaved to produce 34kDa and 14kDa subunits (heavy and light chains, respectively). Cathepsin D exhibits pepsin-like activity and plays a role in protein turnover and in the proteolytic activation of hormones and growth factors. Mutations in this gene play a causal role in neuronal ceroid lipofuscinosis-10 and may be involved in the pathogenesis of several other diseases, including breast cancer and possibly Alzheimer's disease.

Application Notes

Optimal dilution of the CTSD antibody should be determined by the researcher.

Immunogen

Amino acids 104-250 were used as the immunogen of the Cathepsin D antibody.

Storage

Store the Cathepsin D antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).