


Cathepsin D Antibody / CTSD [clone CTSD/2781] (V7476)

Catalog No.	Formulation	Size
V7476-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7476-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7476SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7476IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CTSD/2781
Purity	Protein G affinity chromatography
UniProt	P07339
Localization	Cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Cathepsin D antibody is available for research use only.




Cathepsin D Antibody Renal Cell Carcinoma Immunohistochemistry. IHC testing of FFPE human renal cell carcinoma with Cathepsin D antibody (clone CTSD/2781). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Cathepsin D antibody (clone CTSD/2781). These results demonstrate the foremost specificity of the CTSD/2781 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



Cathepsin D Antibody Liver and MCF7 WB. Western blot testing of human samples with Cathepsin D antibody (clone CTSD/2781).

Description

Cathepsin D is a ubiquitously expressed lysosomal aspartyl protease involved in the normal degradation of proteins. It is synthesized as an inactive 43kDa preprocathepsin D that is cleaved and glycosylated to form a 46kDa procathepsin D and then further cleaved to produce 28kDa and 15kDa subunits (heavy and light chains, respectively). Cathepsin D exhibits pepsin-like activity and plays a role in protein turnover and in the proteolytic activation of hormones and growth factors. Mutations in this gene play a causal role in neuronal ceroid lipofuscinosis-10 and may be involved in the pathogenesis of several other diseases, including breast cancer and possibly Alzheimer's disease.

This antibody can be compared with our [Cathepsin D Antibody / CTSD Antibody](#) (clone CTSD/3276) as a central reference for consistent CTSD detection across multiple assay formats.

Application Notes

Optimal dilution of the Cathepsin D antibody to be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A portion of amino acids 104-250 from the human protein was used as the immunogen for the Cathepsin D antibody.

Storage

Store the Cathepsin D antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

