


CCL13 Antibody / C-C motif chemokine 13 (R31462)

Catalog No.	Formulation	Size
R31462	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
Gene ID	6357
Applications	Western Blot : 0.5-1ug/ml ELISA : 0.1-0.5ug/ml (human protein tested); request BSA free format for coating
Limitations	This CCL13 antibody is available for research use only.



Description

Chemokine (C-C motif) ligand 13 (CCL13), also called SCYL1 and MCP4, is a small cytokine belonging to the CC chemokine family. Its gene is located on human chromosome 17 within a large cluster of other CC chemokines. CCL13 induces chemotaxis in monocytes, eosinophils, T lymphocytes, and basophils by binding cell surface G-protein linked chemokine receptors such as CCR2, CCR3 and CCR5. Activity of this chemokine has been implicated in allergic reactions such as asthma. CCL13 can be induced by the inflammatory cytokines interleukin-1 and TNF-alpha.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the CCL13 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 24-98) was used as the immunogen for this CCL13 antibody.

Storage

After reconstitution, the CCL13 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.