


## CD109 Antibody [clone CD109/9955] (V5675)


Catalog No.	Formulation	Size
V5675-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5675-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5675SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG, kappa
<b>Clone Name</b>	CD109/9955
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q6YHK3
<b>Localization</b>	Cell membrane
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This CD109 antibody is available for research use only.



IHC staining of FFPE human parathyroid tissue with CD109 antibody (clone CD109/9955). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD109 antibody (clone CD109/9955) as confirmation of integrity and purity.

## Description

CD109 modulates negatively TGFB1 signaling in keratinocytes. [UniProt]

## Application Notes

Optimal dilution of the CD109 antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 300-500 from human CD109 protein was used as the immunogen for the CD109 antibody.

## Storage

Aliquot the CD109 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.