
CD11c Antibody ITGAX/1284 / Mouse Monoclonal [clone ITGAX/1284] (V3219)


Catalog No.	Formulation	Size
V3219-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3219-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3219SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	ITGAX/1284
Purity	Protein G affinity chromatography
UniProt	P20702
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CD11c Antibody ITGAX/1284 is available for research use only.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using CD11c Antibody ITGAX/1284. These results demonstrate the foremost specificity of the ITGAX/1284 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



IHC testing of FFPE human tonsil tissue with CD11c antibody (clone ITGAX/1284). Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min.

Description

CD11c Antibody ITGAX/1284 recognizes a protein of 145kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11d (ITGAD), but has a unique alpha chain. CD11c has been shown to play a role in phagocytosis, cell migration, and cytokine production by monocytes/macrophages as well as induction of T-cell proliferation by Langerhans cells. CD11c is expressed prominently on the plasma membranes of monocytes, tissue macrophages, NK cells, and most dendritic cells (DCs). A lower level of expression is also observed on neutrophils as a result of its high level of expression on most DCs. An antibody to CD11c may aid in identification of lesions with histiocytic origin. It may also been used as a marker for hairy cell leukemia in paraffin-embedded tissues.

For a monoclonal antibody with protein microarray specificity validation, see our CD11c Antibody ITGAX/1242 / Integrin Alpha X Antibody page.

Application Notes

Optimal dilution of the CD11c Antibody ITGAX/1284 should be determined by the researcher.

Immunogen

A partial recombinant protein corresponding to amino acids 637-827 from human ITGAX/CD11c was used as the immunogen for the CD11c Antibody ITGAX/1284.

Storage

Store the CD11c Antibody ITGAX/1284 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.