


## CD11c Antibody [clone ITGAX/2507] (V3860)

Catalog No.	Formulation	Size
V3860-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3860-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3860SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


**Bulk quote request**

<b>Availability</b>	1-2 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	ITGAX/2507
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P20702
<b>Localization</b>	Cell surface, cytoplasmic
<b>Applications</b>	ELISA : 1-5ug/ml for coating (order BSA/sodium azide-free format)
<b>Limitations</b>	This CD11c antibody is available for research use only.




IHC testing of FFPE human tonsil with CD11c antibody (clone ITGAX/2507). Required HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

#### Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using CD11c antibody (clone ITGAX/2507). These results demonstrate the foremost specificity of the ITGAX/2507 mAb.

**Z- and S- score:** The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free CD11c antibody (clone ITGAX/2507) as confirmation of integrity and purity.

## Description

Recognizes a protein of 145kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11d (ITGAD), but has a unique alpha chain. CD11c has been shown to play a role in phagocytosis, cell migration, and cytokine production by monocytes/macrophages as well as induction of T-cell proliferation by Langerhans cells. CD11c is expressed prominently on the plasma membranes of monocytes, tissue macrophages, NK cells, and most dendritic cells (DCs). A lower level of expression is also observed on neutrophils as a result of its high level of expression on most DCs. An antibody to CD11c may aid in identification of lesions with histiocytic origin. It may also be used as a marker for hairy cell leukemia in paraffin-embedded tissues.

## Application Notes

Optimal dilution of the CD11c antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein (aa 637-827) was used as the immunogen for the CD11c antibody.

## Storage

Store the CD11c antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).