


CD13 Antibody / ANPEP [clone APN/1464] (V7820)

Catalog No.	Formulation	Size
V7820-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7820-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7820SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	APN/1464
Purity	Protein G affinity chromatography
UniProt	P15144
Applications	ELISA (order BSA-free Format For Coating) :
Limitations	This CD13 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CD13 antibody (clone APN/1464) as confirmation of integrity and purity.

Description

Recognizes an integral membrane glycoprotein of 150kDa, identified as CD13 (also known as aminopeptidase-N). The antibody recognizes an extracellular epitope. The CD13 antigen is present on most cells of myeloid origin including granulocytes, monocytes, mast cells, and GM-progenitor cells. It is also expressed by the majority of AML, CML in

myeloid blast crisis, and in a smaller fraction of lymphoid leukemias. CD13 is absent from normal lymphocytes, platelets and erythrocytes. CD13 is also present on fibroblasts; endothelial cells, epithelial cells from renal proximal tubules and intestinal brush border, bone marrow stromal cells, osteoclasts, and cells lining bile duct canaliculi. CD13 is identical to aminopeptidase N (APN), a prominent membrane-bound metalloprotease present on the surface of intestinal brush border and renal tubules. CD13 plays a role in metabolism of biologically active peptides, in phagocytosis, and in bactericidal/tumoricidal activities. It also serves as a receptor for human coronaviruses (HCV). The lineage-restricted pattern of expression of CD13 within the hemopoietic compartment suggests that it may be important in myeloid cell differentiation.

Application Notes

Optimal dilution of the CD13 antibody should be determined by the researcher.

Immunogen

Recombinant human CD13 protein was used as the immunogen for this CD13 antibody.

Storage

Store the CD13 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).